



## Population structure of the scleractinian coral *Montastraea cavernosa* in southeast Florida

<sup>1</sup>Harbor Branch Oceanographic Institute, Florida Atlantic University, Fort Pierce, Florida 34946

<sup>2</sup>Current address: Cooperative Institute for Marine and Atmospheric Studies (CIMAS), Rosenstiel School of Marine & Atmospheric Science, University of Miami, 4600 Rickenbacker Causeway, Miami, Florida 33149

<sup>3</sup>Florida Fish and Wildlife Conservation Commission, Fort Pierce, Florida 34946

\* Corresponding author email: <jvoss2@fau.edu>

**DL Dodge**<sup>1</sup>  
**MS Studivan**<sup>1,2</sup>  
**RJ Eckert**<sup>1</sup>  
**E Chei**<sup>1</sup>  
**J Beal**<sup>3</sup>  
**JD Voss**<sup>1\*</sup>

---

**ABSTRACT.**—The persistence of scleractinian coral populations on the Florida Reef Tract (FRT) is controlled in part by metapopulation dynamics and larval dispersal. Nine polymorphic microsatellite loci were analyzed to characterize contemporary population structure and gene flow as well as historical migration rates of *Montastraea cavernosa* at five sites off Martin, Palm Beach, and Broward counties in southeast Florida. The sampled populations demonstrated evidence of genetic isolation by distance over a geographic range of 85 km. Population genetic structure was divided into two genetic clusters, northern and southern, with admixture along a latitudinal gradient. Historical migration models indicated likely panmixia throughout all sites sampled, identifying a potential reduction in connectivity among the sampled populations through time. Though *M. cavernosa* populations demonstrated evidence of historical connectivity, contemporary patterns of isolation by distance suggest that effective management will require localized actions to maximize the likelihood of sustaining individual populations in the northern FRT. Given the results of this study, coupled with recent coral mortality events in the region, we recommend regional conservation efforts and management initiatives throughout southeast Florida within a more comprehensive FRT-wide management network.

---

**Handling Editor:** Diego Lirman

Date Submitted: 24 July, 2019.

Date Accepted: 16 December, 2019.

Available Online: 17 December, 2019.

Coral reef ecosystems, and their valuable coastal ecosystem services, have experienced dramatic declines globally since the 1980s (Gardner et al. 2003, Bruno and Selig 2007, Edmunds 2015). Nearshore coral reefs in the Florida Reef Tract (FRT) have suffered severe losses in the past several decades due to increased stressors including land-based sources of pollution, sedimentation, diseases, and thermal stress (Lirman and Fong 2007, Lirman et al. 2011, Ruzicka et al. 2013, Manzello 2015, Precht et al. 2016, Walton et al. 2018). As the third largest barrier reef system in the world, the

FRT extends 595 km between the Dry Tortugas in the southwest and Martin County to the north. Towards the northern limit, this reef system reaches a marginal ecotone between subtropical and warm temperate climate zones (Lugo et al. 1999) with increased environmental variability but diverse marine communities (Beal et al. 2012). The northern FRT (NFRT) in Martin County represents the latitudinal limit in the continental United States for most tropical scleractinian species (Reed 1985). Despite lower coral cover and coral species diversity compared to relatively well-studied reef systems in the Florida Keys and wider tropical western Atlantic, this marginal environment is a significant resource to the local economy and coastal ecosystems (Johns et al. 2004, Collier et al. 2008, Gilliam et al. 2015). Though long-term monitoring and mapping efforts are ongoing (Banks et al. 2007, Walker and Gilliam 2013, Walton et al. 2018), levels of connectivity and gene flow among coral populations in the NFRT are largely unknown.

Understanding patterns of population structure and reef connectivity have become critical in the design and implementation of effective management and restoration strategies. In benthic species, population structure is often driven by larval dispersal (Cowen and Sponaugle 2009), allowing exchange of gametes among populations and possibly an increase in genetic variation required for evolutionary adaptation (van Oppen and Gates 2006). For corals, dispersal potential is dependent on larval characteristics, including spawning strategy (broadcast spawning vs brooding), pelagic larval duration, settlement/survival rates (Cowen and Sponaugle 2009), as well as oceanographic patterns conducive to larval transport. Along the east coast of Florida, the northward-flowing Florida Current, seasonal upwelling, and counter-current eddies are notable influences that can impact the distribution of coral larvae (Limouzy-Paris et al. 1997, Hare and Walsh 2007). Hydrological features and larval characteristics can vary substantially across habitats and coral species, resulting in highly divergent patterns of population connectivity at local and regional scales (Nunes et al. 2011, Bongaerts et al. 2017, Studivan and Voss 2018a, Eckert et al. 2019).

Habitat ranges, larval characteristics, and dispersal potential are difficult to quantify for most marine species including corals (Palumbi 2003, Cowen and Sponaugle 2009). Fortunately, population connectivity across broad temporal and spatial scales can also be investigated through analyses of genomic variations. Population genetic approaches have been commonly used in marine conservation applications, particularly for coral reef ecosystems (Palumbi 2003, Gutiérrez-Rodríguez and Lasker 2004, van Oppen and Gates 2006, Beger et al. 2014, Aswani et al. 2015, Studivan and Voss 2018b). For example, population genetic tools are frequently used in the implementation and design of Marine Protected Areas (MPAs), design of sustainable fisheries, and marine spatial planning through the identification of important metapopulations (Ward et al. 2001, Palumbi 2004). Microsatellite markers, or simple sequence repeats (SSRs), are one common, cost-effective method for investigating population genetic patterns among scleractinian species and geographic locations to answer such conservation-related questions (Miller and Howard 2004, Shearer and Coffroth 2004, Vermeij et al. 2006, Beger et al. 2014, Porto-Hannes et al. 2015).

Previous studies examining population structure across several coral species in the Florida Keys and southern FRT have suggested relatively high connectivity among populations (Baums et al. 2010, Hemond and Vollmer 2010, Serrano et al. 2014, 2016, Manzello et al. 2019). A related study suggested the potential for divergence between populations of *Acropora cervicornis* in the Florida Keys and those in




Figure 1. *Montastraea cavernosa* sampling locations on the northern Florida Reef Tract. Populations were sampled at St Lucie Reef (SLR), Jupiter (JUP), West Palm Beach (WPB), Boynton Beach (BYN), and Fort Lauderdale (FTL). Overlays correspond to reef habitat (shapefile source: Florida Fish and Wildlife Conservation Commission–Fish and Wildlife Research Institute) and the southeast Florida coral reef Ecosystem Conservation Area.

Broward County (Drury et al. 2017), identifying a potential break between northern and southern regions of the FRT. Limited data exists for coral reefs in the NFRT, representing a critical data gap. In conjunction with knowledge of oceanographic patterns influencing larval dispersal, genetic data can inform management strategies that enhance the likelihood of regional survivorship and aid in successful recruitment among downstream populations. Efforts to protect coral reefs along the FRT from further degradation have led to the creation of a southeast Florida coral reef Ecosystem Conservation Area (ECA; FL 2018 House Bill 53; Fig. 1) which extends from St Lucie Inlet to Biscayne National Park. This study was designed to address the population connectivity data gaps in the NFRT and to inform management strategies for the newly established ECA.

*Montastraea cavernosa* is one of the dominant scleractinian coral species in the NFRT and is commonly found throughout the wider tropical western Atlantic as a structurally important reef builder (Nunes et al. 2009). Larval characteristics for *M. cavernosa*, including its relatively large egg size and gonochoric broadcast spawning strategy, may enhance the distance traveled by larvae via oceanographic currents (Szmant 1991). The pelagic larval duration and competency period have not been quantified for this species but are suspected to be relatively long due to observed

Table 1. Site locations and genotyped samples for the northern Florida Reef Tract, with the number of collected samples ( $n$ ) and multi-locus genotypes ( $n_g$ ) used for the analyses. Geographic coordinates given as decimal degrees (WGS84).

| Population (acronym)/Site | Coordinates       | $n$ | $n_g$ | Depth (m) | Collection dates  |
|---------------------------|-------------------|-----|-------|-----------|-------------------|
| St Lucie Reef (SLR)       |                   | 34  | 31    | 5.0       |                   |
| SLR Central               | 27.1317, -80.1340 | 5   | 4     | 2.4       | Sep 2014–Oct 2015 |
| SLR Ledge                 | 27.1214, -80.1275 | 24  | 22    | 9.7       | Jun 2013–Feb 2017 |
| SLR South                 | 27.1119, -80.1255 | 5   | 5     | 3.0       | Dec 2012–Oct 2015 |
| Jupiter (JUP)             |                   | 31  | 30    | 21.0      |                   |
| JUP Ledge                 | 26.9440, -80.0022 | 31  | 30    | 21.0      | Dec 2016          |
| West Palm Beach (WPB)     |                   | 35  | 30    | 17.7      |                   |
| Breaker's Reef            | 26.7179, -80.0187 | 35  | 30    | 17.7      | Jun 2015          |
| Boynton Beach (BYN)       |                   | 33  | 32    | 14.5      |                   |
| SEFL-16                   | 26.5235, -80.0316 | 33  | 32    | 14.5      | Jun 2018          |
| Ft Lauderdale (FTL)       |                   | 31  | 31    | 13.6      |                   |
| BC1                       | 26.1479, -80.0960 | 9   | 9     | 8.2       | Jun 2018          |
| BC2                       | 26.1600, -80.0825 | 13  | 13    | 14.5      | Jun 2018          |
| BC3                       | 26.1586, -80.0774 | 9   | 9     | 18.2      | Jun 2018          |

levels of gene flow among geographically disparate populations. Previous population genetics studies identified broad connectivity across geographic regions in excess of 1000 km (Nunes et al. 2009, Goodbody-Gringley et al. 2012) and vertically among depths in some regions (Brazeau et al. 2013, Serrano et al. 2014, Studivan and Voss 2018a, Eckert et al. 2019). As a common and ecologically important coral in the NFRT, *M. cavernosa* represents an ideal species to examine population structure at multiple spatial scales, allowing inferences to be made across regions and within coral reef habitats of critical ecological and economic importance, such as those found in Florida.

## METHODS

**CORAL SAMPLING.**—Five sites were chosen to assess latitudinal connectivity of *M. cavernosa* along the NFRT: St Lucie Reef (SLR) in Martin County, Jupiter (JUP), West Palm Beach (WPB), Boynton Beach (BYN) in Palm Beach County, and Fort Lauderdale (FTL) in Broward County (Fig. 1). At all locations, coral colonies were selected based on colony size (approximately 20–100 cm diameter), apparent health (no signs of bleaching or disease), and >1 m distance from other sampled colonies to reduce the likelihood of clonal samples. Fragments of coral tissue and skeleton approximately 5 cm<sup>2</sup> in size were collected from the margin of colonies using a hammer and chisel and placed in individual zip-top bags filled with ambient seawater (Table 1). Fragments were transferred to individual tubes and preserved in TRIzol (SLR, JUP, WPB) or molecular grade ethanol (BYN, FTL) and stored at -80 °C until further processing. Differences in sample preservation were due to changes in protocol to reduce user hazards and increase temperature stability of preserved samples; both preservatives yielded sufficient quality genomic DNA for downstream analyses.

**DNA EXTRACTION AND GENOTYPING.**—Frozen TRIzol samples were thawed at room temperature (about 1–2 hrs) and processed with a modified phenol-chloroform

DNA extraction (Chomczynski and Sacchi 2006, Studivan and Eckert 2020). Genomic DNA from ethanol-preserved samples was extracted using a modified Cetyl trimethylammonium bromide (CTAB) extraction protocol (Mieog et al. 2009) as described in Eckert et al. (2019). The resulting DNA pellets were eluted in Tris EDTA (TE) pH 8.0 and incubated at 55 °C for 10 min. DNA extracts were cleaned with the Zymo DNA Clean & Concentrator-5 Kit to remove extraneous proteins and PCR inhibitors. The concentration and quality of all cleaned DNA samples were measured with a NanoDrop 2000 (Thermo Fisher Scientific), and DNA was diluted to a concentration of 10 ng  $\mu\text{l}^{-1}$ . Nine previously-developed microsatellite loci (Serrano et al. 2014) were amplified in all samples with a multiplex PCR using the Qiagen Type-It Microsatellite PCR kit, universal forward primers fluorescently-labeled with NED, VIC, or 6-FAM, and reverse primers (Blacket et al. 2012; Online Table S1). PCR thermal cycling consisted of the following: 95 °C for 5 min (initial denaturation), 30 (ethanol samples) or 35 (TRIzol samples) cycles of 30 s at 95 °C (denaturation), 60 °C for 90 s (annealing), 72 °C for 30 s (extension), and 60 °C for 30 min (final extension). Additional PCR cycles were necessary for TRIzol samples due to variable initial DNA quality.

Amplified alleles were visualized on a 2% agarose gel to verify amplification success and to select dilution strength based on band intensity. PCR products were diluted and sized with an Applied Biosystems 3130 XL (500 ROX size standard) at Florida Atlantic University's Harbor Branch Oceanographic Institute. Resulting electropherograms were analyzed and alleles were scored with GeneMapper v3.7 (Applied Biosystems). Samples with low or unidentifiable allele peaks at individual loci were reamplified to correct for incomplete data. Samples were run a mean of 1.3 times per locus on the genetic analyzer to ensure sufficient coverage across all loci. After re-amplification attempts, samples missing data at more than three loci were removed from further analyses.

**GENETIC AND STATISTICAL ANALYSES.**—Multilocus genotypes (MLGs) and violations of Hardy-Weinberg Equilibrium (HWE) across loci were identified in GenAIEx v6.5 (Peakall and Smouse 2012). Genetic diversity metrics, including population differentiation, fixation index ( $F_{ST}$ ), and Nei's genetic distance ( $D$ ) were calculated in GenAIEx. Private alleles per locus within populations were determined using the *R* package *diveRsity* (Keenan et al. 2013). Linkage disequilibrium (LD) assumptions were tested in Arlequin v3.5 (Excoffier and Lischer 2010). All  $P$  values for HWE, LD, and  $F_{ST}$  were corrected for false discovery rate (FDR; Benjamini and Hochberg 1995). Violations of HWE and LD assumptions were tested for significant impact of null alleles, where null allele-corrected  $F_{ST}$  values were calculated using FreeNA (Chapuis and Estoup 2007), then compared to raw  $F_{ST}$  values using correlation analyses across populations and loci. Loci were examined for HWE and LD violations and patterns of allelic richness across populations to determine if any loci needed to be removed prior to downstream analyses. An analysis of molecular variance (AMOVA) was calculated among individuals and populations using GenAIEx with 9999 permutations. A principal coordinates analysis (PCoA) visualized genetic distance between populations using Nei's genetic distance, and a Mantel isolation by distance test assessed the relationship between genetic and geographic distance in GenAIEx (9999 permutations) using over-water distances calculated from site GPS coordinates.

Structure v2.3.4 (Pritchard et al. 2000) was used to analyze genetic structure and identify populations. Samples were assigned to  $K$  populations in a probabilistic manner until HWE and linkage equilibrium was achieved, assuming that loci were unlinked and able to freely recombine. Using an admixture model, simulations were run with  $10^6$  Markov Chain-Monte Carlo (MCMC) replications after a 1000 burn-in period. The number of postulated populations ( $K$ ) was tested between 1 and 8 with 10 replicate tests per value of  $K$ . Structure output results were analyzed using the web-based Structure Harvester (Earl and VonHoldt 2012) to determine the most likely value of  $K$  using the Evanno method (Evanno et al. 2005). A combined structure plot of all replicate simulations for the highest likelihood  $K$  was created to represent individual membership to genetic clusters using CLUMPP v1.1.2 (Jakobsson and Rosenberg 2007) and Distruct v1.1 (Rosenberg 2004).

Historical migration rates of ancestral *M. cavernosa* populations (approximately  $a4N_e$  generations) among sites were estimated with Migrate-n v3.6.11 (Beerli 2006, Beerli and Palczewski 2010) using Bayesian inference to determine the maximum likelihood of various genetic migration models. Generation times and mutation rates for microsatellite loci are unknown for most coral species (*but see* Devlin-Durante et al. 2016). Because *M. cavernosa* is relatively long-lived and reproductively viable at an early age (Szmant 1991, Soong 1993), the migration estimates provided by Migrate-n represent patterns over hundreds to thousands of years. Four possible migration models were tested: (1) symmetrical migration across all sites; (2) unidirectional migration from southern to northern sites; (3) unidirectional migration from northern to southern sites; and (4) panmixia. All models allowed stepwise migration, where upstream populations could contribute migrants to all downstream populations (and vice versa for the symmetrical migration model). All migration models were run with the following parameters: 20 replicates, long-inc 100, long-sample 15,000, burn-in 20,000, and 4 heated chains (1, 1.5, 3,  $10^5$ ). Prior distributions for mutation-scaled population size ( $\Theta$ ) and mutation-scaled immigration rate ( $M$ ) were set from 0 to 100 and 0 to 1000, respectively. The most likely migration scenario was determined by ranking of Bezier log marginal likelihoods [ $\ln(m)$ ] for each model using the thermodynamic integration method described in Beerli and Palczewski (2010). To assess whether the genetic divergence of the Fort Lauderdale population from all other populations may have been driving historical migration results, the model scenarios were repeated excluding Fort Lauderdale from the dataset.

## RESULTS

Of the nine microsatellite markers amplified in this study, eight consistently met HWE assumptions across populations (Online Table S2). No violations of LD assumptions were observed across loci or populations. In addition to HWE violations in three out of five sampled populations, locus MC4 demonstrated abnormally high numbers of alleles ( $N_a$ ) and allelic richness ( $A_r$ ; Online Table S2). Allele bins within locus MC4 were also close to one another, compromising the ability to definitively make allele calls. MC4 was therefore removed from all further analyses. Subsequent analyses using the remaining dataset identified no clonal MLGs within the populations, allowing all amplified samples to be used for statistical assessments of genetic connectivity. Null allele-corrected  $F_{ST}$  values were found to be tightly correlated with raw  $F_{ST}$  values across loci and populations ( $R^2 = 0.9926$  and  $R^2 = 0.9976$ , respectively); therefore, raw values were reported.




Figure 2. Principal coordinates analysis (PCoA) of Nei's genetic distance ( $D_A$ ) among pairwise comparisons of populations, explaining 78.57% of the total genetic variation (PCo 1: 56.97%; PCo 2: 21.60%). Color of points corresponds to the dominant genetic cluster identified in Structure analysis ( $K = 2$ ; Fig. 5).

Allelic diversity was similar among populations, with an overall mean range of allelic richness ( $A_r$ ) between 3.54 and 19.50. Fort Lauderdale demonstrated the lowest richness ( $A_r = 2.79$ ), while Jupiter had the greatest ( $A_r = 22.75$ ). Expected heterozygosity ( $H_e$ ) was also similar among populations, with Fort Lauderdale once again having the lowest ( $H_e = 0.16$ ) and Jupiter having the highest ( $H_e = 0.95$ ). The global fixation index ( $F_{ST}$ ) was 0.017 across all populations. An analysis of molecular variance (AMOVA) indicated that 6% of genetic variation was within populations ( $df = 149$ ,  $SS = 436.932$ ,  $P = 0.0001$ ), while only 2% of observed variation was among populations ( $df = 4$ ,  $SS = 23.888$ ,  $P = 0.0001$ ). PCoA demonstrated clustering of northernmost sites St Lucie Reef and Jupiter, indicating high genetic similarity of these populations (Fig. 2). Genetic distance among sites reflected site geography, with West Palm Beach and Boynton Beach being intermediately distinct, and the southernmost site Fort Lauderdale being the most dissimilar population. Pairwise population  $F_{ST}$  values reinforced the PCoA results, indicating that Fort Lauderdale was significantly differentiated from all other sites and supporting genetic isolation of sites by distance




Figure 3. Pairwise  $F_{ST}$  values among sampled geographic populations. Bolded values indicate significantly differentiated populations and color intensity reflects level of genetic differentiation from  $F_{ST}$  values. Alpha was set at 0.05.




Figure 4. Mantel isolation by distance test relating Nei's genetic distance ( $D$ ) to geographic distance among pairwise comparisons of populations. Geographic isolation explained 39.46% of observed genotypic variation among populations (Mantel test;  $P < 0.033$ , 999 permutations).

(Fig. 3). The Mantel test revealed a significant correlation between geographic distance and Nei's genetic distance ( $R^2 = 0.3946$ ,  $P = 0.033$ ; Fig. 4). Geographic distances among pairwise comparisons of sites ranged from 21.65 to 84.57 km.

A Bayesian population clustering model in Structure identified the most likely number of genetic clusters ( $K$ ) among populations to be  $K = 2$  through both log model likelihood [ $L(K)$ ] and the Evanno method (Online Table S3, Figs. 5 and Online S1). Northernmost sites St Lucie Reef and Jupiter exhibited similar genetic structure and were derived predominantly from a single genetic cluster represented by dark blue in Figure 5. The southernmost site Fort Lauderdale was dominated by a separate genetic cluster represented by light blue. Intermediate sites West Palm Beach and Boynton Beach showed admixture from both populations; however, both were primarily represented by the same genetic cluster observed at the northern sites (dark blue). Assessment of historical population migration using Migrate-n indicated panmixia among all sampled populations. Model likelihood for panmictic migration was much higher than for other tested models simulating stepwise symmetric, south-north, and north-south migration, with near-zero probabilities for the latter models (Table




Figure 5. Genetic structure of *Montastraea cavernosa* populations in the northern Florida Reef Tract as predicted by Bayesian inference in Structure. Each column represents an individual *M. cavernosa* colony from the five geographic populations (x-axis). The relative height of each color (dark blue, light blue) represents the probability of membership to the genetic clusters ( $K = 2$ ).



Table 2. Comparison of Bezier log marginal likelihoods [ $\ln(mL)$ ] for migration models (symmetric migration, migration from south to north, migration from north to south, and panmixia) across all populations, and following the removal of Fort Lauderdale (FTL). Models were ranked according to likelihood values to calculate probabilities and rank. Mutation-scaled population size ( $\Theta$ ) included for the most probable model (with 95% CI).

| Model           | $\ln(mL)$ | Probability | Rank | $\Theta$ (95% CI)  |
|-----------------|-----------|-------------|------|--------------------|
| All populations |           |             |      |                    |
| Symmetric       | -871,446  | 0           | 4    |                    |
| South-North     | -475,069  | 0           | 3    |                    |
| North-South     | -288,996  | 0           | 2    |                    |
| Panmixia        | -26,436   | 1           | 1    | 12.50 (4.87–10.27) |
| FTL removed     |           |             |      |                    |
| Symmetric       | -413,402  | 0           | 4    |                    |
| South-North     | -78,427   | 0           | 3    |                    |
| North-South     | -58,799   | 0           | 2    |                    |
| Panmixia        | -20,073   | 1           | 1    | 8.49 (3.40–7.80)   |

2). Repeated model simulations using a dataset excluding Fort Lauderdale demonstrated that panmixia was once again the most likely migration scenario, with all other migration models showing near-zero probabilities. Estimated population sizes for both likely models are given in Table 2.

## DISCUSSION

This study provides evidence that populations of an important reef-building coral, *M. cavernosa*, in the northern Florida Reef Tract were historically connected. However, examination of contemporary population structure and gene flow among *M. cavernosa* populations in the NFRT identified patterns of genetic differentiation. *Montastraea cavernosa* exhibits relatively high levels of horizontal gene flow across its range and particularly throughout the Florida Keys, Gulf of Mexico, Belize, Barbados, Bermuda, Jamaica, and Panama (Nunes et al. 2009, Goodbody-Gringley et al. 2012, Serrano et al. 2014, Studivan and Voss 2018a, Eckert et al. 2019). Certain larval characteristics are thought to increase the ability of pelagic larvae to survive and therefore travel long distances, thus increasing population connectivity throughout a larger region (Hare and Walsh 2007). The relatively large egg size of *M. cavernosa*, broadcast spawning reproductive mode, and pelagic larval duration period may enable broad transport of larvae across geographically-isolated areas (Szmant 1991, Acosta and Zea 1997, Jones et al. 2009), provided oceanographic patterns allow transport among reefs. While pelagic larval duration and competency periods have not been quantified for *M. cavernosa*, it is not suspected that larval characteristics reduce this species' dispersal potential, given the evidence of connectivity observed over ranges exceeding 1000 km (Nunes et al. 2009, 2011, Goodbody-Gringley et al. 2012, Studivan and Voss 2018a).

Oceanographic patterns are likely the main driver of larval transport and gene flow among populations in Florida. The predominant Florida Current, which is fed by the consistently strong (typically  $>1 \text{ m s}^{-1}$ ) Gulf of Mexico Loop Current through the Straits of Florida (Oey et al. 2005), runs nearly the entire length of the FRT and has the potential to transport larvae from the southern to the northern extent of the FRT (Limouzy-Paris et al. 1997, Hare and Walsh 2007, Sponaugle and Cowen

2019). Nearshore habitats also experience localized upwelling and counter-current eddies that deviate shoreward from the Florida Current, which have been shown to influence larval dispersal patterns (Limouzy-Paris et al. 1997, Sponaugle et al. 2005). Given that hydrological features in the region have been predicted to be relatively consistent and supportive of reef growth through the Holocene beginning around 12,000 ya (Banks et al. 2008), it is not surprising that we observed historical panmixia in *M. cavernosa* populations across a linear distance of approximately 85 km.

While *M. cavernosa* populations in the NFRT demonstrate historical connectivity based on the Migrate-n results, there is additional evidence of relatively recent population differentiation based on contemporary population structure and gene flow. Multiple analyses demonstrate isolation by distance for *M. cavernosa* in the study region (Fig. 4). Additionally, evaluation of population structure identified two dominant genetic clusters across the range of sampled sites (Fig. 5). Northernmost sites St Lucie Reef and Jupiter were nearly indistinguishable from one another while the southernmost site, Fort Lauderdale, was distinct. Admixture at intermediate sites West Palm Beach and Boynton Beach indicated variable influences of both genetic clusters across a latitudinal gradient. It is important to note, however, that observed differences in population structure among sites appeared to be largely driven by pairwise differences between Fort Lauderdale and all other sampled sites (Fig. 3). Given these observations, it appears that the dark blue genetic cluster common across the NFRT demonstrates higher genetic diversity than the light blue cluster predominantly found in Fort Lauderdale. For example, Jupiter was shown to have the highest allelic richness (also the highest membership to the dark blue cluster), while Fort Lauderdale had the lowest.

The results of these complementary analyses suggest that population sources are different between sites in Martin and Palm Beach counties (St Lucie Reef, Jupiter, West Palm Beach, and Boynton Beach) and those in Broward County (Fort Lauderdale). To our knowledge, no studies have examined coral population structure across all ranges of the FRT, but previous population genetics analyses of *A. cervicornis*, a broadcast spawning species, suggest a further divergence between populations in the Florida Keys and those in Broward County (Drury et al. 2017). It is therefore possible that coral populations may be split into zones along the FRT, with variable patterns of gene flow and genetic diversity within and among regions. The observed trends in this study may be further explained by variation in oceanographic patterns that occur at the northern extent of the FRT. Following the Atlantic continental shelf, the Florida Current diverges from the Florida coastline at the Bahamas Fault Zone where coastal morphology changes (Walker 2012). The two northernmost sites sampled for this study, St Lucie Reef and Jupiter, are located north of the Bahamas Fault Zone, potentially reducing northward transport of larvae from southern populations. Subregional oceanographic features including eddies between the Florida Current and nearshore habitats may be sufficient to allow gene flow over time, albeit at a lower rate of success with increasing latitude. This is perhaps best evidenced by the decreasing level of admixture between genetic clusters from southern to northern sites (Fig. 5).

The more recent population divergence observed in the NFRT may be indicative of habitat degradation and declines in *M. cavernosa* abundance along the FRT (Beal et al. 2012, Walton et al. 2018) with associated reductions in coral reproduction and recruitment. Coral cover and reef diversity have declined in recent decades in part

due to macroalgal phase shifts (Palandro et al. 2008, Burman et al. 2012, Ruzicka et al. 2013), severe mortality events following thermal stress (Lirman et al. 2011), and disease outbreaks (Precht et al. 2016). Likewise, reduced coral recruitment has been observed elsewhere in the tropical western Atlantic (Hughes 1994, Mumby et al. 2007, Mumby 2009). Coral mortality events may constitute genetic bottlenecks, where winnowed coral populations can lead to decreased genetic diversity and gene flow. Spawning and recruitment have not been examined for most of the NFRT (*but see* Miller et al. 2000, Lirman and Fong 2007). Colony fate-tracking of *M. cavernosa* at St Lucie Reef revealed no observed spawning in the past decade and no histological evidence of gamete production (Beal et al. 2012, Klepac et al. 2015). It may also be possible that exogenous *M. cavernosa* recruits still reach the NFRT but do not survive into adulthood or become reproductive members of the population.

**IMPLICATIONS FOR MANAGEMENT.**—Our results suggest that contemporary *M. cavernosa* populations in the NFRT do not appear to be a substantial contributor to regional larval supply. Instead, it is more likely that populations are considered sinks, with additional divergence between the northern (St Lucie Reef, Jupiter, West Palm Beach, and Boynton Beach) and southern (Fort Lauderdale) regions of the NFRT. There is a need to quantitatively track recruitment among populations on the FRT, allowing the identification of important source populations and perhaps the prioritization of regions in need of conservation initiatives, especially with respect to ongoing coral nursery and gene banking programs. Therefore, this study's results provide support for regional conservation through a network of small-scale management initiatives throughout southeast Florida within an FRT-wide comprehensive network. Resolving patterns of genetic structure over small spatial scales is required to make informed local management guidelines, especially with diverse reef habitats in the NFRT where coral density is low and reproduction (both sexual and asexual) is reduced. Such factors decrease the ability and likelihood of these reefs to recover from mortality events, leading to an increased need to sustain and protect present resources. Informed management of coral reefs in the southeast Florida coral reef ECA will require (1) additional sampling to fill in the gaps for the dominant coral, *M. cavernosa*, in southern Broward and Miami-Dade counties, and (2) similar approaches for additional targeted coral and fish species.

An increasing number of studies are examining multiple aspects of connectivity within regions, particularly in the comparison of genetic and biophysical connectivity (Galindo et al. 2006, Davies et al. 2017, Garavelli et al. 2018, Studivan and Voss 2018b). The FRT, with its dynamic current patterns and variable reef diversity across latitudes, would benefit greatly from studies of this nature. There is a critical need for detailed sampling and observation of distinct gradients in the region, ideally with molecular markers capable of higher resolution, such as single nucleotide polymorphism (SNP) loci (Drury et al. 2017, Manzello et al. 2019). This need is particularly acute given continued coral population declines resulting from the recent stony coral tissue loss disease outbreak along the FRT (Precht et al. 2016, Walton et al. 2018).

#### ACKNOWLEDGMENTS

We are grateful to the St Lucie Inlet Preserve State Park, Martin County, Florida Fish and Wildlife Conservation Commission, and the Florida Department of Environmental

Protection Coral Reef Conservation Program for their collaboration and support of this research. We thank A Shatters, J Polinski, A Alker, P Gardner, A Sturm, I Combs, C Luck, J Nelson, and M Roy for outstanding diving and boat support. T Ferrer and G O’Corry-Crowe at FAU HBOI provided Applied Biosystems 3130 XL sequencing services. This is contribution 2264 from FAU’s Harbor Branch Oceanographic Institute. This project was supported by Florida Sea Grant R/LR-B-70 to J Voss and J Beal, NOAA Coral Reef Conservation Program grant NA16NOS4820052 to J Voss, EPA grant X700D66717 to J Voss and J Beal, PADI Foundation grant 21853 to D Dodge, a Women Divers Hall of Fame scholarship to D Dodge, an Indian River Lagoon Graduate Research Fellowship to D Dodge, and a donation from the River Branch Foundation. Corals were collected under Florida Fish and Wildlife Conservation Commission authority and Florida State Park Service permit 09231515.

#### LITERATURE CITED

- Acosta A, Zea S. 1997. Sexual reproduction of the reef coral *Montastrea cavernosa* (Scleractinia: Faviidae) in the Santa Marta area, Caribbean coast of Colombia. *Mar Biol.* 128:141–148. <https://doi.org/10.1007/s002270050077>
- Aswani S, Mumby P, Baker AC, Christie P, McCook LJ, Steneck RS, Richmond RH. 2015. Scientific frontiers in the management of coral reefs. *Front Mar Sci.* 2:1–13. <https://doi.org/10.3389/fmars.2015.00050>
- Banks KW, Riegl BM, Richards VP, Walker BK, Helmle KP, Jordan LKB, Phipps J, Shivji MS, Spieler RE, Dodge RE. 2008. The reef tract of continental Southeast Florida (Miami-Dade, Broward and Palm Beach Counties, USA). *Coral Reefs of the USA*. Dordrecht: Springer Netherlands. p. 175–220.
- Banks KW, Riegl BM, Shinn EA, Piller WE, Dodge RE. 2007. Geomorphology of the Southeast Florida continental reef tract (Miami-Dade, Broward, and Palm Beach Counties, USA). *Coral Reefs.* 26:617–633. <https://doi.org/10.1007/s00338-007-0231-0>
- Baums IB, Johnson ME, Devlin-Durante MK, Miller MW. 2010. Host population genetic structure and zooxanthellae diversity of two reef-building coral species along the Florida Reef Tract and wider Caribbean. *Coral Reefs.* 29:835–842. <https://doi.org/10.1007/s00338-010-0645-y>
- Beal J, Voss JD, Edge S, Cohen L. 2012. Assessment of coral stressors on St Lucie Reef: Florida’s northernmost coral reef. Final report for State Wildlife Grant. USFWS T-91-1, Fort Pierce, Florida. 54 p.
- Berli P. 2006. Comparison of Bayesian and maximum-likelihood inference of population genetic parameters. *Bioinformatics.* 22:341–345. <https://doi.org/10.1093/bioinformatics/bti803>
- Berli P, Palczewski M. 2010. Unified framework to evaluate panmixia and migration direction among multiple sampling locations. *Genetics.* 185:313–326. <https://doi.org/10.1534/genetics.109.112532>
- Beger M, Selkoe KA, Treml E, Barber PH, Von Der Heyden S, Crandall ED, Toonen RJ, Riginos C. 2014. Evolving coral reef conservation with genetic information. *Bull Mar Sci.* 90:159–185. <https://doi.org/10.5343/bms.2012.1106>
- Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc B.* 57:289–300. <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x>
- Blacket MJ, Robin C, Good RT, Lee SF, Miller AD. 2012. Universal primers for fluorescent labelling of PCR fragments-an efficient and cost-effective approach to genotyping by fluorescence. *Mol Ecol Resour.* 12:456–463. <https://doi.org/10.1111/j.1755-0998.2011.03104.x>
- Bongaerts P, Riginos C, Brunner R, Englebort N, Smith SR, Hoegh-Guldberg O. 2017. Deep reefs are not universal refuges: reseeded potential varies among coral species. *Sci Adv.* 3:e1602373. <https://doi.org/10.1126/sciadv.1602373>

- Brazeau DA, Lesser MP, Slattery M. 2013. Genetic structure in the coral, *Montastraea cavernosa*: assessing genetic differentiation among and within mesophotic reefs. PLOS ONE. 8:e65845. <https://doi.org/10.1371/journal.pone.0065845>
- Bruno JF, Selig ER. 2007. Regional decline of coral cover in the Indo-Pacific: timing, extent, and subregional comparisons. PLOS ONE. 2:e711. <https://doi.org/10.1371/journal.pone.0000711>
- Burman SG, Aronson RB, van Woesik R. 2012. Biotic homogenization of coral assemblages along the Florida reef tract. Mar Ecol Prog Ser. 467:89–96. <https://doi.org/10.3354/meps09950>
- Chapuis M-P, Estoup A. 2007. Microsatellite null alleles and estimation of population differentiation. Mol Biol Evol. 24:621–631. <https://doi.org/10.1093/molbev/msl191>
- Chomczynski P, Sacchi N. 2006. The single-step method of RNA isolation by acid guanidinium thiocyanate–phenol–chloroform extraction: twenty-something years on. Nat Protoc. 1:581–585. <https://doi.org/10.1038/nprot.2006.83>
- Collier C, Ruzicka R, Banks K, Barbieri L, Beal J, Bingham D, Bohnsack JA, Brooke S, Craig N, Dodge RE, et al. 2008. The state of coral reef ecosystems of Southeast Florida. In: Waddell J, Clarke A, editors. The state of coral reef ecosystems of the United States and Pacific Freely Associated States: 2008 NOAA Technical Memorandum NOS NCCOS 73. NOAA/NCCOS. Silver Spring, Maryland: Center for Coastal Monitoring and Assessment's Biogeography Team. p. 131–160.
- Cowen RK, Sponaugle S. 2009. Larval dispersal and marine population connectivity. Annu Rev Mar Sci. 1:443–466. <https://doi.org/10.1146/annurev.marine.010908.163757>
- Davies SW, Strader ME, Kool JT, Kenkel CD, Matz MV. 2017. Modeled differences of coral life-history traits influence the refugium potential of a remote Caribbean reef. Coral Reefs. 36:913–925. <https://doi.org/10.1007/s00338-017-1583-8>
- Devlin-Durante MK, Miller MW, Precht WF, Baums IB, Carne L, Smith TB, Banaszak AT, Greer L, Irwin A, Fogarty ND, et al. 2016. How old are you? Genet age estimates in a clonal animal. Mol Ecol. 25:5628–5646. <https://doi.org/10.1111/mec.13865>
- Drury C, Schopmeyer S, Goergen E, Bartels E, Nedimyer K, Johnson M, Maxwell K, Galvan V, Manfrino C, Lirman D. 2017. Genomic patterns in *Acropora cervicornis* show extensive population structure and variable genetic diversity. Ecol Evol. 7:6188–6200. <https://doi.org/10.1002/ece3.3184>
- Earl DA, VonHoldt BM. 2012. STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. Conserv Genet Resour. 4:359–361. <https://doi.org/10.1007/s12686-011-9548-7>
- Eckert RJ, Studivan MS, Voss JD. 2019. Populations of the coral species *Montastraea cavernosa* on the Belize Barrier Reef lack vertical connectivity. Sci Rep. 9:7200. <https://doi.org/10.1038/s41598-019-43479-x>
- Edmunds PJ. 2015. A quarter-century demographic analysis of the Caribbean coral, *Orbicella annularis*, and projections of population size over the next century. Limnol Oceanogr. 60:840–855. <https://doi.org/10.1002/lno.10075>
- Evanno G, Regnaut S, Goudet J. 2005. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. Mol Ecol. 14:2611–2620. <https://doi.org/10.1111/j.1365-294X.2005.02553.x>
- Excoffier L, Lischer HEL. 2010. Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. Mol Ecol Resour. 10:564–567. <https://doi.org/10.1111/j.1755-0998.2010.02847.x>
- Galindo HM, Olson DB, Palumbi SR. 2006. Seascape genetics: a coupled oceanographic-genetic model predicts population structure of Caribbean corals. Curr Biol. 16:1622–1626. <https://doi.org/10.1016/j.cub.2006.06.052>
- Garavelli L, Studivan MS, Voss JD, Kuba A, Figueiredo J, Chérubin LM. 2018. Assessment of mesophotic coral ecosystem connectivity for proposed expansion of a marine sanctuary in the

- northwest Gulf of Mexico: larval dynamics. *Front Mar Sci.* 5:174. <https://doi.org/10.3389/fmars.2018.00174>
- Gardner TA, Côté IM, Gill JA, Grant A, Watkinson AR. 2003. Long-term region-wide declines in Caribbean corals. *Science.* 301:958–960. <https://doi.org/10.1126/science.1086050>
- Gilliam DS, Walton CJ, Brinkhuis V, Ruzicka R, Colella M. 2015. Southeast Florida coral reef evaluation and monitoring project 2014. Year 12 Final Report, Miami Beach, Florida. 43 p.
- Goodbody-Gringley G, Woollacott RM, Giribet G. 2012. Population structure and connectivity in the Atlantic scleractinian coral *Montastraea cavernosa* (Linnaeus, 1767). *Mar Ecol.* 33:32–48. <https://doi.org/10.1111/j.1439-0485.2011.00452.x>
- Gutiérrez-Rodríguez C, Lasker HR. 2004. Microsatellite variation reveals high levels of genetic variability and population structure in the gorgonian coral *Pseudopterogorgia elisabethae* across the Bahamas. *Mol Ecol.* 13:2211–2221. <https://doi.org/10.1111/j.1365-294X.2004.02247.x>
- Hare JA, Walsh HJ. 2007. Planktonic linkages among marine protected areas on the south Florida and southeast United States continental shelves. *Can J Fish Aquat Sci.* 64:1234–1247. <https://doi.org/10.1139/f07-089>
- Hemond EM, Vollmer SV. 2010. Genetic diversity and connectivity in the threatened staghorn coral (*Acropora cervicornis*) in Florida. *PLOS ONE.* 5:e8652. <https://doi.org/10.1371/journal.pone.0008652>
- Hughes TP. 1994. Catastrophes, phase shifts, and large-scale degradation of a Caribbean coral reef. *Science.* 265:1547–1551. <https://doi.org/10.1126/science.265.5178.1547>
- Jakobsson M, Rosenberg NA. 2007. CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics.* 23:1801–1806. <https://doi.org/10.1093/bioinformatics/btm233>
- Johns GM, Milon J, Sayers D. 2004. Socioeconomic study of reefs in Martin County, Florida. Report prepared for Martin County, Florida by Hazen and Sawyer, PC, Hollywood, Florida. 120 p.
- Jones GP, Almany GR, Russ GR, Sale PF, Steneck RS, van Oppen MJH, Willis BL. 2009. Larval retention and connectivity among populations of corals and reef fishes: history, advances and challenges. *Coral Reefs.* 28:307–325. <https://doi.org/10.1007/s00338-009-0469-9>
- Keenan K, McGinnity P, Cross TF, Crozier WW, Prodöhl PA. 2013. diveRsity: an R package for the estimation of population genetics parameters and their associated errors. *Methods Ecol Evol.* 4:782–788. <https://doi.org/10.1111/2041-210X.12067>
- Klepac CN, Beal J, Kenkel CD, Sproles A, Polinski JM, Williams MA, Matz MV, Voss JD. 2015. Seasonal stability of coral-*Symbiodinium* associations in the subtropical coral habitat of St Lucie Reef, Florida. *Mar Ecol Prog Ser.* 532:137–151. <https://doi.org/10.3354/meps11369>
- Limouzy-Paris CB, Graber HC, Jones DL, Röpke AW, Richards WJ. 1997. Translocation of larval coral reef fishes via sub-mesoscale spin-off eddies from the Florida current. *Bull Mar Sci.* 60:966–983.
- Lirman D, Fong P. 2007. Is proximity to land-based sources of coral stressors an appropriate measure of risk to coral reefs? An example from the Florida Reef Tract. *Mar Pollut Bull.* 54:779–791. <https://doi.org/10.1016/j.marpolbul.2006.12.014>
- Lirman D, Schopmeyer S, Manzello D, Gramer LJ, Precht WF, Muller-Karger F, Banks K, Barnes B, Bartels E, Bourque A. 2011. Severe 2010 cold-water event caused unprecedented mortality to corals of the Florida Reef Tract and reversed previous survivorship patterns. *PLOS ONE.* 6:e23047. <https://doi.org/10.1371/journal.pone.0023047>
- Lugo AE, Brown SB, Dodson R, Smith TS, Shugart HH. 1999. The Holdridge life zones of the conterminous United States in relation to ecosystem mapping. *J Biogeogr.* 26:1025–1038. <https://doi.org/10.1046/j.1365-2699.1999.00329.x>
- Manzello DP. 2015. Rapid recent warming of coral reefs in the Florida Keys. *Sci Rep.* 5:16762. <https://doi.org/10.1038/srep16762>
- Manzello DP, Matz MV, Enochs IC, Valentino L, Carlton RD, Kolodziej G, Serrano X, Towle EK, Jankulak M. 2019. Role of host genetics and heat-tolerant algal symbionts in sustaining

- populations of the endangered coral *Orbicella faveolata* in the Florida Keys with ocean warming. *Glob Change Biol.* 25:1016–1031. <https://doi.org/10.1111/gcb.14545>
- Mieog JC, van Oppen MJH, Berkelmans R, Stam WT, Olsen JL. 2009. Quantification of algal endosymbionts (*Symbiodinium*) in coral tissue using real-time PCR. *Mol Ecol Resour.* 9:74–82. <https://doi.org/10.1111/j.1755-0998.2008.02222.x>
- Miller KJ, Howard CG. 2004. Isolation of microsatellites from two species of scleractinian coral. *Mol Ecol Notes.* 4:11–13. <https://doi.org/10.1046/j.1471-8286.2003.00555.x>
- Miller MW, Weil E, Szmant AM. 2000. Coral recruitment and juvenile mortality as structuring factors for reef benthic communities in Biscayne National Park, USA. *Coral Reefs.* 19:115–123. <https://doi.org/10.1007/s003380000079>
- Mumby PJ. 2009. Phase shifts and the stability of macroalgal communities on Caribbean coral reefs. *Coral Reefs.* 28:761–773. <https://doi.org/10.1007/s00338-009-0506-8>
- Mumby PJ, Hastings A, Edwards HJ. 2007. Thresholds and the resilience of Caribbean coral reefs. *Nature.* 450:98–101. <https://doi.org/10.1038/nature06252>
- Nunes FLD, Norris RD, Knowlton N. 2009. Implications of isolation and low genetic diversity in peripheral populations of an amphi-Atlantic coral. *Mol Ecol.* 18:4283–4297. <https://doi.org/10.1111/j.1365-294X.2009.04347.x>
- Nunes FLD, Norris RD, Knowlton N. 2011. Long distance dispersal and connectivity in amphi-Atlantic corals at regional and basin scales. *PLOS ONE.* 6:e22298. <https://doi.org/10.1371/journal.pone.0022298>
- Oey L, Ezer T, Lee H. 2005. Loop Current, rings and related circulation in the Gulf of Mexico: a review of numerical models and future challenges. *In: Sturges W, Lugo-Fernandez A, editors. Circulation in the Gulf of Mexico: observations and models.* Washington DC: American Geophysical Union. p. 31–56.
- Palandro DA, Hu C, Hallock P, Müller-Karger FE, Dustan P, Callahan MK, Kranenburg C, Beaver CR. 2008. Quantification of two decades of shallow-water coral reef habitat decline in the Florida Keys National Marine Sanctuary using Landsat data (1984–2002). *Remote Sens Environ.* 112:3388–3399. <https://doi.org/10.1016/j.rse.2008.02.015>
- Palumbi SR. 2003. Population genetics, demographic connectivity, and the design of marine reserves. *Ecol Appl.* 13:146–158. [https://doi.org/10.1890/1051-0761\(2003\)013\[0146:PGD CAT\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2003)013[0146:PGD CAT]2.0.CO;2)
- Palumbi SR. 2004. Marine reserves and ocean neighborhoods: the spatial scale of marine populations and their management. *Annu Rev Environ Resour.* 29:31–68. <https://doi.org/10.1146/annurev.energy.29.062403.102254>
- Peakall R, Smouse PE. 2012. GenA1Ex 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics.* 28:2537–2539. <https://doi.org/10.1093/bioinformatics/bts460>
- Porto-Hannes I, Zubillaga AL, Shearer TL, Bastidas C, Salazar C, Coffroth MA, Szmant AM. 2015. Population structure of the corals *Orbicella faveolata* and *Acropora palmata* in the Mesoamerican Barrier Reef System with comparisons over Caribbean basin-wide spatial scale. *Mar Biol.* 162:81–98. <https://doi.org/10.1007/s00227-014-2560-1>
- Precht WF, Gintert BE, Robbart ML, Fura R, van Woesik R. 2016. Unprecedented disease related coral mortality in southeastern Florida. *Sci Rep.* 6:31374. <https://doi.org/10.1038/srep31374>
- Pritchard JK, Stephens M, Donnelly P. 2000. Inference of population structure using multilocus genotype data. *Genetics.* 155:945–959.
- Reed JK. 1985. Deepest distribution of Atlantic hermatypic corals discovered in the Bahamas. *Proceedings of the 5th International Coral Reef Congress, Tahiti.* 6:249–254.
- Rosenberg NA. 2004. DISTRUCT: a program for the graphical display of population structure. *Mol Ecol Notes.* 4:137–138. <https://doi.org/10.1046/j.1471-8286.2003.00566.x>
- Ruzicka RR, Colella MA, Porter JW, Morrison JM, Kidney JA, Brinkhuis V, Lunz KS, MacAulay KA, Bartlett LA, Meyers MK, et al. 2013. Temporal changes in benthic assemblages on

- Florida Keys reefs 11 years after the 1997/1998 El Niño. *Mar Ecol Prog Ser.* 489:125–141. <https://doi.org/10.3354/meps10427>
- Schuelke M. 2000. An economic method for the fluorescent labeling of PCR fragments. *Nat Biotechnol.* 18:233–234. <https://doi.org/10.1038/72708>
- Serrano XM, Baums IB, O'Reilly K, Smith TB, Jones RJ, Shearer TL, Nunes FLD, Baker AC. 2014. Geographic differences in vertical connectivity in the Caribbean coral *Montastraea cavernosa* despite high levels of horizontal connectivity at shallow depths. *Mol Ecol.* 23:4226–4240. <https://doi.org/10.1111/mec.12861>
- Serrano XM, Baums IB, Smith TB, Jones RJ, Shearer TL, Baker AC. 2016. Long-distance dispersal and vertical gene flow in the Caribbean brooding coral *Porites astreoides*. *Sci Rep.* 6:21619. <https://doi.org/10.1038/srep21619>
- Shearer TL, Coffroth MA. 2004. Isolation of microsatellite loci from the scleractinian corals, *Montastraea cavernosa* and *Porites astreoides*. *Mol Ecol Notes.* 4:435–437. <https://doi.org/10.1111/j.1471-8286.2004.00653.x>
- Soong K. 1993. Colony size as a species character in massive reef corals. *Coral Reefs.* 12:77–83. <https://doi.org/10.1007/BF00302106>
- Sponaugle S, Cowen RK. 2019. Coral ecosystem connectivity between Pulley Ridge and the Florida Keys. *In: Loya Y, Puglise KA, Bridge TCL, editors. Mesophotic coral ecosystems, coral reefs of the world 12.* Switzerland: Springer Nature. p. 897–907.
- Sponaugle S, Lee T, Kourafalou V, Pinkard D. 2005. Florida Current frontal eddies and the settlement of coral reef fishes. *Limnol Oceanogr.* 50:1033–1048. <https://doi.org/10.4319/lo.2005.50.4.1033>
- Studivan MS, Eckert RJ. 2020. mstudiva/Mcav-microsats: Final *M. cavernosa* microsatellite pipeline (Version 1.0.0). Zenodo. <http://doi.org/10.5281/zenodo.3728179>
- Studivan MS, Voss JD. 2018a. Population connectivity among shallow and mesophotic *Montastraea cavernosa* corals in the Gulf of Mexico identifies potential for refugia. *Coral Reefs.* 37:1183–1196. <https://doi.org/10.1007/s00338-018-1733-7>
- Studivan MS, Voss JD. 2018b. Assessment of mesophotic coral ecosystem connectivity for proposed expansion of a marine sanctuary in the Northwest Gulf of Mexico: population genetics. *Front Mar Sci.* 5:152. <https://doi.org/10.3389/fmars.2018.00152>
- Szmant AM. 1991. Sexual reproduction by the Caribbean reef corals *Montastrea annularis* and *M. cavernosa*. *Mar Ecol Prog Ser.* 74:13–25. <https://doi.org/10.3354/meps074013>
- van Oppen MJH, Gates RD. 2006. Conservation genetics and the resilience of reef-building corals. *Mol Ecol.* 15:3863–3883. <https://doi.org/10.1111/j.1365-294X.2006.03026.x>
- Vermeij MJA, Fogarty ND, Miller MW. 2006. Pelagic conditions affect larval behavior, survival, and settlement patterns in the Caribbean coral *Montastraea faveolata*. *Mar Ecol Prog Ser.* 310:119–128. <https://doi.org/10.3354/meps310119>
- Walker BK. 2012. Spatial analyses of benthic habitats to define coral reef ecosystem regions and potential biogeographic boundaries along a latitudinal gradient. *PLOS ONE.* 7:e30466. <https://doi.org/10.1371/journal.pone.0030466>
- Walker BK, Gilliam DS. 2013. Determining the extent and characterizing coral reef habitats of the northern latitudes of the Florida Reef Tract (Martin County). *PLOS ONE.* 8:e80439. <https://doi.org/10.1371/journal.pone.0080439>
- Walton CJ, Hayes NK, Gilliam DS. 2018. Impacts of a regional, multi-year, multi-species coral disease outbreak in Southeast Florida. *Front Mar Sci.* 5:323. <https://doi.org/10.3389/fmars.2018.00323>
- Ward TJ, Heinemann D, Evans N. 2001. The role of marine reserves as fisheries management tools. A review of concepts, evidence and international experience. Canberra, Australia: Bureau of Rural Sciences. 192 p.

